



**AIDS
2012**

**XIX INTERNATIONAL AIDS
CONFERENCE JULY 22 - 27
WASHINGTON DC USA**



Friday 27th July 2012

Contents

- [How long until a cure?](#)
- [One year's preventive therapy reduces the risk of TB in people on HIV treatment](#)
- [HIV and TB](#)
- [Keeping people in HIV care](#)
- [Retaining children in HIV care](#)
- [Access to treatment](#)
- [HIV and the criminal law](#)
- [HIV and hepatitis C](#)
- [Hormonal contraception and HIV risk](#)
- [The global village](#)
- [Support our work](#)



How long until a cure?



The panel at the HIV cure press conference. © IAS/Deborah W. Campos - Commercialimage.net

Some of the latest HIV cure research has been presented at International AIDS Conference this week.

Experts also met for a [cure workshop prior to the conference](#), where they launched a global scientific strategy *Towards an HIV Cure*.

Research is looking at a range of different approaches to a possible cure, including:

- flushing out and destroying HIV lying dormant in 'reservoirs' in the body.
- stem cell treatment (like that which cured the '[Berlin Patient](#)')
- starting HIV treatment very soon after infection – an approach that would only work for a small proportion of people with HIV.

It's likely any successful cure process will involve a combination of approaches.

Promising results in some of the cure studies raise their own ethical questions, as people who are doing well on successful HIV treatment would have to stop to see if a functional cure has been achieved. An ethics working group has been set up to address these issues.

Steven Deeks, co-chair with Francois Barré-Sinoussi of the IAS HIV Cure working group, said: "The barriers to a cure are far greater than barriers to antiretroviral therapy [in the late 1980s]... Unless we get very lucky this is going to take well over a decade."

"The field is moving fast", said Sharon Lewin from Monash University in Melbourne. "We certainly don't have a cure currently, but we have a better understanding of what we need to do."

Related links

[View full news story on aidsmap](#)

[View abstract THAA0101 on the conference website](#)

[View abstract THAA0103 on the conference website](#)

One year's preventive therapy reduces the risk of TB in people on HIV treatment


Related links

[View full news report on aidsmap](#)

[View the abstract on the conference website](#)

[View the webcast of the presentation on the conference website.](#)

Keeping people in HIV care



Images from the presentation of Dr Rachel Baggaley of the World Health Organisation.

Community-based support increases the likelihood that HIV-positive people will be retained in care, research conducted in South Africa shows.

Less than a third of people who start HIV therapy in southern Africa stay in HIV care. This is of serious concern. It means that the majority of people with HIV are not receiving the health benefits that come from specialist treatment and care. A high drop-out rate from care also seriously undermines the use of HIV treatment as prevention.

A speaker from the World Health Organization (WHO) presented findings from research in more than 20 countries about why people fall out of care at some point in the 'cascade' of HIV care (that is, the different stages of care from first being tested to being on effective treatment). Reasons include the fear of stigma, denial about their condition, anxiety, poor links with the care available, inadequate clinic facilities and problems with travel.

But a South African study showed that providing community-based adherence support increased the chances that people starting HIV treatment would stay in care.

A new healthcare role of 'patient advocates' was introduced in 2004. These workers help support adherence and also provide counselling and psychosocial support.

Only 6% of people who had an advocate dropped out of care, compared to 10% of individuals who did not receive this kind of support.

Separate research showed that contact tracing could help reduce drop-out rates among children after starting HIV treatment.

Following the introduction of active contact tracing, rates of loss-to-follow-up were reduced from 22.7% to 8.5%.

The WHO report recommends the involvement of lay health workers to help ensure people move

from one stage of the care 'cascade' to the next, and therefore stay in care.

In partnership with UNICEF



Related links

[View full news report on aidsmap](#)

[View information on the session, including links to the abstracts and slides from the presentations, on the conference website](#)

Retaining children in HIV care



Rene Ekpini of UNICEF. © IAS/Steve Shapiro - Commercialimage.net

On the same theme of keeping people in care, [the Washington conference was told that three-quarters of children in need of HIV therapy are not receiving treatment, an unacceptable situation.](#)

However, a variety of schemes show that this situation can be remedied.

One, [implemented in Malawi](#), includes the provision of lifelong HIV treatment to all HIV-positive pregnant women, regardless of their CD4 cell count.

Another intervention in Zimbabwe has increased rates of diagnoses among children.

A number of practical measures – often with the local community playing a pivotal role – were also improving the retention of children in care.

These included:

- Use of community volunteers to accompany children to clinic appointments.
- Patient advocacy.
- Provision of transport vouchers.

In partnership with UNICEF

Related links

[View full news report on aidsmap](#)

[View information on the satellite session, with links to slides, on the conference website](#)

[View abstract THAE0101 on the conference website](#)

[View the slides from the presentation on the conference website](#)

Access to treatment



Protester at the Say it loud! march in Washington. Image by Greta Hughson/aidsmap.com

Patents and intellectual property rights are still restricting access to antiretroviral therapy in middle- and low-income countries, the Washington conference was told.

The roll-out of antiretroviral therapy in poorer countries has been made possible in part due to the development of cheaper, generic formulations of a number of key [anti-HIV drugs](#).

However, the maintenance of intellectual property rights means that medications needed for second- and third-line treatment remain prohibitively expensive. The conference also heard of the especially high cost of treatment for some middle-income countries.

Speakers at the session recommended that efforts were made to challenge patent applications to ensure access to treatment takes priority.

Related links

[View full news story on aidsmap](#)

[View information on the session, including links to the abstracts and slides from the presentations, on the conference website](#)

HIV and the criminal law



Edwin J Bernard of the HIV Justice Network. © IAS/Ryan Rayburn - Commercialimage.net

Draconian laws on HIV transmission and exposure were the focus of a session at the International AIDS Conference.

Delegates heard that courts often dismissed evidence regarding the excellent prognosis of patients on HIV therapy and the impact of HIV treatment on the risk of transmission.

People with HIV are being imprisoned after sexual encounters when they did not disclose their status, even when no transmission occurred. In some instances, people had been prosecuted even though the type of sex they had engaged in involved no actual risk of HIV transmission.

More encouragingly, the session was also told that lobbying could lead to changes in the law.

For instance, Denmark suspended its tough HIV-specific laws after being presented with scientific evidence about the life expectancy of people on effective HIV therapy and the impact of treatment as prevention.

Related links

[View full news report on aidsmap](#)

[View information on the session, including links to the abstracts, slides and webcasts from the presentations, on the conference website](#)

HIV and hepatitis C



Vincent Lo Re of the University of Pennsylvania. Image ©Liz Highleyman / hivandhepatitis.com

People co-infected with HIV and hepatitis C continue to have a higher risk of serious liver disease than those who only have hepatitis C.

Researchers in the US compared rates of liver disease and liver-related death between co-infected and hepatitis C-monoinfected people.

Importantly, the co-infected participants were on HIV treatment, which has previously been shown to slow the progression of liver disease.

Co-infected people were at approximately twice the risk of developing decompensated liver disease and 69% more likely to progress to liver cancer.

An undetectable HIV viral load reduced the risk of liver disease, but even with HIV suppression outcomes were still poorer in co-infected people compared to those who only had hepatitis C.

Related links

[View full news report on aidsmap](#)

[View the abstract on the conference website](#)

[View the slides from the presentation on the conference website](#)

Hormonal contraception and HIV risk



Contradictory evidence concerning the use of hormonal contraception and its possible impact on the risk of infection with HIV was presented to the International AIDS Conference.

Research published last year showed that women who used hormonal contraception had an increased risk of infection with HIV and were also more likely to transmit the virus.

A re-analysis of the results of this study confirmed the association between hormonal contraception and an increased risk of acquiring HIV.

The association between the use of hormonal contraception and HIV risk remained significant when taking into account rates of unprotected sex.

However, a meta-analysis of studies looking at the risk of HIV and the use of hormonal and non-hormonal forms of contraception failed to find any definitive evidence that hormonal contraception increased the chances of infection with HIV.

The point was made that access to reliable contraception is important for women, and that any possible HIV risk has to be balanced with the availability of effective contraception.

Related links

[View full news report on aidsmap](#)

[View a complete listing of the session presentations, with links to abstracts and slides, on the conference website](#)

The global village



Images by Greta Hughson/aidsmap.com

The **19th International AIDS Conference (AIDS 2012)** is not just the sum of its presentations and posters. Alongside the conference sessions, the exhibition halls and the global village have also been a hive of activity this week.

Join **aidsmap's** Greta Hughson on a tour of the **global village**.

Related links

[The Global Village](#)

Support our work



As a charity we rely on donations to continue our work and are so grateful for every gift we receive, no matter how big or small. Every single penny is put towards helping people with HIV, and those who support and care for them, access the vital information they need.

We believe passionately that independent, clear and evidence-based information lies at the heart of empowering people to make decisions about their health and live longer, healthier, happier lives.

Read about **how your support makes a difference** and if you can feel you can support our work


with a donation, you can do so online at www.aidsmap.com/donate.

Thank you.

"I was diagnosed with HIV in early December at the age of 24... I initially couldn't bring myself to talk to even my closest friends or family about the diagnosis and aidsmap.com has helped me beyond belief and helped me to be able to talk to others in person."

Related links

 www.aidsmap.com/donate



Two other official partners are providing coverage and analysis online, so you can have the fullest picture of the conference. **Clinical Care Options (CCO)**, will be providing audio highlights, capsule summaries and downloadable slidesets, while the **Kaiser Family Foundation** are providing webcasting from conference sessions.



Connect with NAM on Facebook: Keep up to date with all the exciting projects, latest achievements and new developments that are going on in the world of NAM.



Follow NAM on twitter for links to hot off the press news stories from our editors covering key developments and conferences as they happen. Our news feed is linked to www.twitter.com/aidsmap_news and we also tweet from www.twitter.com/aidsmap.



Follow all the conference news by **subscribing to our RSS feeds**.

AIDS 2012 conference coverage in partnership with:



NAM is partnering with UNICEF to deliver the AIDS 2012 bulletins, which have also been made possible thanks to support from Bristol-Myers Squibb and Vestergaard Frandsen. NAM's wider conference news reporting services have been supported by Abbott, Boehringer Ingelheim, Janssen, Roche and ViiV Healthcare.



NAM is an award-winning, community-based organisation, which works from the UK. We deliver

reliable and accurate HIV information across the world to HIV-positive people and to the professionals who treat, support and care for them.

Make a donation, make a difference at www.aidsmap.com/donate

For more details, please contact NAM:

tel: +44 (0)20 3242 0820

fax: +44 (0)20 3242 0839

email: info@nam.org.uk

web: www.aidsmap.com

NAM Publications

Registered office: 77a Tradescant Road, London, SW8 1XJ

Company limited by guarantee. Registered in England & Wales, number: 2707596

Registered charity, number: 1011220

To unsubscribe please visit: <http://www.aidsmap.com/page/1492854/>